Physical Variables written by CTIPe simulations

The coordinate system (geographic) consists of

- Geographic longitude Lon with positive vector components meaning eastward,
- Geographic latitude Lat from -90 at the south pole to 90 at the north pole with **positive** being **northward**,
- pressure level **IP** or height **H** in km.

Vector (arrow) plots of the velocities only make sense as:

- vertical cuts (meridional or at constant-latitude) if UseHeight is selected,
- **synoptic maps** of velocity vectors (over local time and latitude) if plotted at **constant height** (not constant pressure level IP).

The basic plasma and electrodynamic field variables in 3D are:

- Neutral mass density **rho** in [kg/m³].
- **H** (height) in [km] corresponding to pressure level number **IP**The height of a pressure level varies spatially and with time. Heights covered start at about 80 km (**IP**=0) and reach a few hundred km above ground (the maximum found for **IP**=14, the top layer, is typically between 450 km and 1000 km).

The height can be used as an alternative 3rd coordinate for plotting.

- Particle number density N in [m⁻³] with species identifier (after the " "):
 - o e: electrons,
 - o **O**: atomic oxygen,
 - N2: nitrogen molecules,
 - o **O2**: oxygen molecules,
 - o NO: nitric oxide,
 - NO+: nitric oxide ion,
 - o N2+: molecular nitrogen ion,
 - o **O2+**: molecular oxygen ion,
 - N+: atomic nitrogen ion,
 - o **O**+: atomic oxygen ion,
 - H+: atomic hydrogen ion.
- Neutral gas temperature T_n in [K].
- Mean molecular mass **Rmt** in [amu].
- Hall and Pedersen conductivities sigma_H, sigma_P in [mho/m].
- Neutral gas velocity Vn in [m/s] with its three components
 - Vn Lat (meridional; CTIP name "V x"),
 - Vn Lon (zonal, longitudinal; CTIP name "V y") and
 - Vn IP (vertical, radial; CTIP name "V z").

Pjoule: joule heating in [J/(kg·sec)]

Prad: radiation heating/cooling in [J/(kg·sec)]

· Electric field

E140_theta: latitudinal component of electric field at 140 km in [V/m]

E140 lambda: longitudinal component of electric field at 140 km in [V/m]

E300 theta: latitudinal component of electric field at 300 km in [V/m]

E300 lambda: longitudinal component of electric field at 300 km in [V/m]

Height-integrated quantities in 3D data

available at each position in local time and latitude (obtained from 3D CTIP variables above)

- NmF2: maximum electron density N_e in [m⁻³] in the vertical profile,
- HmF2: height in [km] of the maximum of N e (see NmF2),
- TEC: integrated total electron content in [TECU=10¹⁶ electrons/m²] in the altitude range of 80 km 2000 km.

Height-integrated quantities

- **Wjoule**: Joule heating [mW/m²],
- Win: Energy flux [mW/m²],
- En_avg: Mean particle energy [keV],

Energy deposition rates (in GW)

- P tot: auroral energy input over both the northern and southern hemispheres
- **P_euv,N**: extreme ultraviolet solar radiation ($\lambda < 102.7$ nm) integrated over northern hemisphere
- **P_euv,S**: extreme ultraviolet solar radiation ($\lambda < 102.7$ nm) integrated over southern hemisphere
- **P_uv,N**: far ultraviolet solar radiation (102.7 nm $< \lambda <$ 200 nm) integrated over northern hemisphere
- **P_uv,S**: far ultraviolet solar radiation (102.7 nm $< \lambda <$ 200 nm) integrated over southern hemisphere
- P J.E.N: sum of Joule heating and kinetic energy dissipation in northern hemisphere
- P J.E.S: sum of Joule heating and kinetic energy dissipation in northern hemisphere
- P Joule, N: joule heating integrated over the northern hemisphere
- **P_Joule,S**: joule heating integrated over the northern hemisphere
- **P_kin,N**: kinetic energy in the northern hemisphere
- **P_kin,S**: kinetic energy in the southern hemisphere
- P kin: kinetic energy in both the southern and northern hemispheres

Changes in output parameters from geometric quiet condition (Kn. 2): rd(output parameter)